

Joseph V. Candela, MD
Vijay Goli, MD, FACS
Victor E. Grigoriev, MD, FACS
Steven B. Kurtz, MD

William B. Steinkohl, MD, FACS
O. Alex Lesani, MD
William R. Wise, MD
Jeffrey M. Zapinsky, MD, FACS



Diplomates of the
American Board of Urology

POST OPERATIVE INSTRUCTIONS FOR STONE SURGERY

IMPORTANT PLEASE READ COMPLETELY

1. After your stone surgery, you may continue to experience pain either due to the procedure or to an indwelling tube left in the kidney and bladder to promote healing. Continue taking your pain pills (Lortab, Percocet, or Darvocet N-100) as needed. If this doesn't control the pain, please contact the doctor immediately.
2. If there was a tube left in your kidney and bladder, burning and frequent urination is common. Take your Pyridium (Phenazopyridine) or Proceid EC as directed to help with these symptoms. This medicine normally turns the urine orange or blue. This tube will be taken out in the office as soon as possible. Once it is removed, the above mentioned symptoms usually resolve immediately.
3. Drink plenty of fluids. Any type of fluid (non-alcoholic) is acceptable. If you experience significant nausea or are unable to take enough fluids, call the doctor.
4. Take your antibiotics as directed.
5. Someone from our staff will call you the following business day during office hours to check on your condition and schedule your follow-up appointment. If you haven't been contacted within this time frame, please contact our office as soon as possible. You may be asked to get x-rays prior to your visit with the doctor. Please bring these x-rays with you so the doctor can assess the results of your surgery.

NORTHWEST

7500 Smoke Ranch
#200 - Las Vegas, NV 89128
TELEPHONE: (702) 233-0727

SOUTHWEST

7150 W. Sunset Rd.
#201A - Las Vegas, NV 89113
FAX: (702) 233-4799